

AD-776 059

PLUME CHEMISTRY MODELS

SUMMARY OF A PLUME CHEMISTRY WORKSHOP

PHYSICAL DYNAMICS, INCORPORATED

PREPARED FOR

ADVANCED RESEARCH PROJECTS AGENCY

AIR FORCE CAMBRIDGE RESEARCH LABORATORIES

1 APRIL 1973

DISTRIBUTED BY:

**NTIS**

National Technical Information Service  
U. S. DEPARTMENT OF COMMERCE

AD776059

AFCRL-TR-73-0697

PD-73-036

PLUME CHEMISTRY MODELS  
SUMMARY OF A PLUME CHEMISTRY WORKSHOP

by

Edward R. Fisher


PHYSICAL DYNAMICS INC.  
P.O. Box 604  
College Park Station  
Detroit, Michigan 48221

Contract No. F19628-72-C-0006  
Project No. 8692

Scientific Report No. 3

April 1, 1973

Contract Monitor: Alva T. Stair  
Optical Physics Laboratory



Approved for public release; distribution unlimited.

Reproduced by  
NATIONAL TECHNICAL  
INFORMATION SERVICE  
U S Department of Commerce  
Springfield VA 22151

Sponsored by  
Defense Advanced Research Projects Agency  
ARPA Order No. 1856

Monitored by  
Air Force Cambridge Research Laboratories  
Air Force Systems Command  
United States Air Force  
Bedford, Massachusetts 01730

//

ARPA Order No. 1856

Program Code No. 1E40

Contractor: Physical Dynamics, Inc.

Effective Date of Contract: 1 July 1972

Contract No. F19628-72-C-0006

Principal Investigator and Phone No.

Mr. Frederick P. Boynton/(415)-848-3063

AFCRL Project Scientist and Phone No.

Dr. Alva T. Stair/(617)-861-4911

Contract Expiration Date: 31 December 1973

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
D C	Def Section <input type="checkbox"/>
UNCLASSIFIED	<input type="checkbox"/>
JUSTIFICATION .....	
BY .....	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	Avail. and/or SPECIAL
A	

Qualified requestors may obtain additional copies from the Defense Documentation Center. All others should apply to the National Technical Information Service.

UNCLASSIFIED

Security Classification

## DOCUMENT CONTROL DATA - R &amp; D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Physical Dynamics, Inc. P.O. Box 604, College Park Station Detroit, Michigan 48221		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
3. REPORT TITLE  PLUME CHEMISTRY MODELS, SUMMARY OF A PLUME CHEMISTRY WORKSHOP		2b. GROUP	
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Scientific. Interim.			
5. AUTHOR(S) (First name, middle initial, last name)  Edward R. Fisher			
6. REPORT DATE April 1, 1973		7a. TOTAL NO. OF PAGES 27	7b. NO. OF REFS 5
8a. CONTRACT OR GRANT NO. ARPA Order No. 1856 F19628-72-C-0006		9a. ORIGINATOR'S REPORT NUMBER(S) PD 73-036, Scientific Report #3	
b. PROJECT NO. 8692		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) AFCRL-TR-73-0697	
c. DOD ELEMENT 62301D			
d. DOD SUBELEMENT n/a			
10. DISTRIBUTION STATEMENT  A - Approved for public release; distribution unlimited.			
11. SUPPLEMENTARY NOTES This research was sponsored by the Defense Advanced Research Projects Agency.		12. SPONSORING MILITARY ACTIVITY Air Force Cambridge Research Labs (OP) L.G. Hanscom Field Bedford, Mass. 01730	
13. ABSTRACT  Two plume chemistry models are presented as a result of a workshop meeting in January, 1973. The first model characterizes the chemi-excitation and relaxation processes appropriate to an O/H/OH/H <sub>2</sub> /O <sub>2</sub> /H <sub>2</sub> O system, while the second model characterizes the relaxation processes in a mixture of CO <sub>2</sub> , H <sub>2</sub> O, H <sub>2</sub> , CO, N <sub>2</sub> , O and H. These chemistry models are presented for baseline predictions on plume behavior.			

DDC  
RECEIVED  
MAR 27 1974  
B

DD FORM 1473  
NOV 66REPLACES DD FORM 1473, 1 JAN 64, WHICH IS  
OBSOLETE FOR ARMY USE.

UNCLASSIFIED

Security Classification

UNCLASSIFIED

Security Classification

14.	KEY WORDS		LINK A		LINK B		LINK C	
			ROLE	WT	ROLE	WT	ROLE	WT
	Plume Chemistry Models Relaxation Model APOLLO Model							

iA

Security Classification

## TABLE OF CONTENTS

	<u>page</u>
ABSTRACT	iii
INTRODUCTION	1
CHEMICAL MODELS	3
Table I.    APOLLO Chemistry Model	6
References	9
Table II.  Vibrational Relaxation Model	11
References	16
Table III. Radiative Lifetimes	18
SUMMARY	19
BIBLIOGRAPHY	20
APPENDIX I	
Attendance List to Plume Chemistry Workshop	21

## INTRODUCTION

As the plume modeling capabilities continue to develop under the ARPA Plume Physics Program, a reliable set of "baseline" chemical mechanisms characterizing various fuel/atmospheric environments becomes increasingly important. This importance stems not only from providing direct comparisons between various flow field calculations using the same chemical models but also in establishing a basis upon which to judge improvements in reaction mechanisms and rate coefficients as new experimental and theoretical results become available. In this report, two chemical mechanisms with accompanying rate coefficients are presented: a chemi-excitation and vibrational relaxation mechanism for the APOLLO vehicle ( $H_2/O_2$  System), and a general vibrational relaxation mechanism for a system consisting of  $H_2O$ ,  $CO_2$ ,  $CO$ ,  $N_2$ ,  $O$ ,  $H$  and  $OH$  species. These reaction mechanisms are the outgrowth of a chemistry workshop held at Physical Dynamics, Inc., Berkeley, California on January 12, 1973. A list of attendees to this workshop are given in Appendix I. It should be noted that the collective input for all attending is represented in these two reaction mechanisms, particularly with respect to the magnitude of the proposed rate coefficients.

It will be clear upon reviewing the references accompanying the reaction mechanisms that many rate coefficients are very poorly known particularly for reactions involving atomic and radical species. Experiments underway at Wayne State University, AVCO and CALSPAN under the Plume Physics Program are expected to improve these estimates in the coming months.


In proposing these baseline chemical mechanisms, it should be noted that an attempt was made to include all types of chemical reactions. Thus, reactions with relatively high activation energies generally limited in importance to high temperature regions of the plume are included as well as fast reactions involving reactive intermediates such as OH.

The mechanisms presented in this report should complement the "lumped parameter" hydrocarbon and amine combustion mechanisms recently proposed [Kummler, Fisher, Boynton (1972)] as well as the recent review of vibrational cross section data appropriate to rarified plume environments [Fisher (1972)].

## CHEMICAL MODELS

The chemical reactions listed in Table I for the APOLLO system consist of both chemi-excitation and vibrational relaxation processes. To date there is no direct information on the rate coefficients for the chemi-excitation reactions [listed as reactions 1) through 18)] except for the overall reactions as reviewed in the Leed's reports [Baulch (1969)]. The rate coefficients for the reactions involving excited states have been estimated following a method of Boynton (1972) which most likely consists of an upper limit to these excited state reactions. The approach taken in these estimates is to assume that vibrational energy in the reactants [e.g.,  $H_2(v=1)$  as shown in reaction 3)] removes the experimentally observed overall activation energy as compiled in the Leed's reviews. Thus, the high temperature region (above about 1500°K) is dominated by the excited state reaction where  $H_2(v=1)$  is present in relatively large concentrations while the low temperature reaction is dominated by the  $H_2(v=0)$  reaction. By assuming a Boltzmann population of  $H_2(v=1)$  at any given temperature, the additive contribution of reactions 1) and 3) are made to fit the measured overall reaction rate.

The reverse rate coefficients for reactions involving excited states have been calculated from the appropriate equilibrium constant as taken from the Leeds's review together with Boltzmann factors to account for the excited state populations. For example, the equilibrium constant for the reaction set



is related to the equilibrium constant for the overall reaction



as given in the Leeds reports by

$$K_{56} = \frac{e^{-E_1/KT}}{e^{-E_2/KT}} \frac{Q_2}{Q_1} K_{\text{Leeds}}$$

where

$E_1$  = vibrational spacing of OH

$E_2$  = vibrational spacing of  $\text{H}_2$

$Q_1$  = vibrational partition function of OH

$Q_2$  = vibrational partition function of  $\text{H}_2$

and  $K_{\text{Leeds}}$  = thermodynamic equilibrium constant for the overall reaction.

The reverse rate coefficients for chemical excitation reactions producing  $\text{H}_2\text{O}$  [i.e. reactions 7), 9), 11) and 13)] were calculated from the forward rate coefficients assuming that the  $\text{H}_2\text{O}$  is formed in the vibrational ground state. Note also that the energy transfer reactions [i.e. reactions 14) through 47) and Table II] are only listed in the forward direction. The reverse rate coefficients can be calculated from the Boltzmann factor involving the energy defect (energy available to the translational and rotational modes).

Table III lists the best available Einstein A Coefficients for incorporation into plume chemistry models. Of course the prediction of radiated energy in any given wavelength region depends not only on the total energy available from a given vibrational state as given by the A Coefficient, but also on the spectral shape of that band (i.e. the rotational distribution). A more extensive discussion of this point can be found elsewhere [Boynton, Fisher, Kummeler, Thomson (1973)].

TABLE I  
APOLLO CHEMISTRY MODEL

<u>Reaction</u>	<u>Reference</u>	<u>Rate Constant</u> (cc/molecule-sec)
1) $O + H_2 \rightarrow OH + H$	(a,b)	$1.65^{-13} \exp(-2500/T)$
2) $H + OH \rightarrow O + H_2$	(c)	$6.9^{-14} \exp(-1425/T)$
3) $O + H_2(v=1) \rightarrow OH + H$	(d)	$1.3^{-10} \alpha \ (\alpha = 0.9, 0.5)$
4) $H + OH \rightarrow O + H_2(v=1)$	(c)	$5.5^{-11} \alpha \exp(-4925/T)$
5) $O + H_2(v=1) \rightarrow OH(v=1) + H$	(d)	$1.3^{-10} (1 - \alpha)$
6) $H + OH(v=1) \rightarrow O + H_2(v=1)$	(c)	$5.5^{-11} (1 - \alpha) \exp(215/T)$
7) $OH + H_2 \rightarrow H_2O + H$	(d)	$3.8^{-11} \exp(-2600/T)$
8) $H + H_2O \rightarrow OH + H_2$	(c)	$1.5^{-10} \exp(-10075/T)$
9) $OH + H_2(v=1) \rightarrow H_2O + H$	(e)	$1.5^{-10}$
10) $H + H_2O \rightarrow OH + H_2(v=1)$	(c)	$5.8^{-10} \exp(-13475/T)$
11) $OH(v=1) + H_2 \rightarrow H_2O + H$	(e)	$1.5^{-11}$
12) $H + H_2O \rightarrow OH(v=1) + H_2$	(c)	$5.8^{-11} \exp(-12475/T)$
13) $OH + OH \rightarrow H_2O + O$	(a)	$9.6^{-12} \exp(-390/T)$
14) $O + H_2O \rightarrow OH + OH$	(a)	$9.5^{-11} \exp(-9000/T)$
15) $O + OH \rightarrow O_2 + H$	(f)	$4^{-11}$
16) $H + O_2 \rightarrow O + OH$	(c)	$6.9^{-10} \exp(-8400/T)$
17) $O + OH(v=1) \rightarrow O_2 + H$	(f)	$4^{-11}$
18) $H + O_2 \rightarrow O + OH(v=1)$	(c)	$6.9^{-10} \exp(-13400/T)$

# ENERGY TRANSFER REACTIONS\*

## Vibration-Translation Processes:

19) $\text{H}_2(v=1) + \text{H} \rightarrow \text{H}_2 + \text{H}$	(g)	$3.2^{-11} \exp(-1400/T)$
20) $\text{H}_2(v=1) + \text{O} \rightarrow \text{H}_2 + \text{O}$	(e)	$3.0^{-13} \sqrt{T}$
21) $\text{H}_2(v=1) + \text{H}_2 \rightarrow \text{H}_2 + \text{H}_2$	(h,i)	$1.5^{-7} \exp(-139/T^{1/3})$
22) $\text{H}_2(v=1) + \text{H}_2\text{O} \rightarrow \text{H}_2 + \text{H}_2\text{O}$	(e,j)	$1.5^{-7} \exp(-139/T^{1/3})$
23) $\text{N}_2(v=1) + \text{H} \rightarrow \text{N}_2 + \text{H}$	(u)	$1.9^{-6} \exp(-164/T^{1/3})$
24) $\text{N}_2(v=1) + \text{O} \rightarrow \text{N}_2 + \text{O}$	(k,l,m)	$1.2^{-13} \exp(-23/T^{1/3})$
25) $\text{N}_2(v=1) + \text{H}_2 \rightarrow \text{N}_2 + \text{H}_2$	(n)	$3.9^{-8} \exp(-164/T^{1/3})$
26) $\text{N}_2(v=1) + \text{H}_2\text{O} \rightarrow \text{N}_2 + \text{H}_2\text{O}$	(e,j)	$3.9^{-8} \exp(-164/T^{1/3})$
27) $\text{OH}(v=1) + \text{H} \rightarrow \text{OH} + \text{H}$	(e)	$3^{-11}$
28) $\text{OH}(v=1) + \text{O} \rightarrow \text{OH} + \text{O}$	(e)	$3^{-11}$
29) $\text{OH}(v=1) + \text{H}_2 \rightarrow \text{OH} + \text{H}_2$	(e)	$1^{-11}$
30) $\text{OH}(v=1) + \text{H}_2\text{O} \rightarrow \text{OH} + \text{H}_2\text{O}$	(e)	$3^{-12}$
31) $\text{H}_2\text{O}(010) + \text{H} \rightarrow \text{H}_2\text{O}(000) + \text{H}$	(e)	$1^{-11}$
32) $\text{H}_2\text{O}(010) + \text{O} \rightarrow \text{H}_2\text{O}(000) + \text{O}$	(e)	$1^{-11}$
33) $\text{H}_2\text{O}(010) + \text{H}_2 \rightarrow \text{H}_2\text{O}(000) + \text{H}_2$	(e,o)	$1^{-11}$
34) $\text{H}_2\text{O}(010) + \text{H}_2\text{O} \rightarrow \text{H}_2\text{O}(000) + \text{H}_2\text{O}$	(p)	$1^{-11}$
35) $\text{H}_2\text{O}(020) + \text{H} \rightarrow \text{H}_2\text{O}(010) + \text{H}$	(t)	$2^{-11}$
36) $\text{H}_2\text{O}(020) + \text{O} \rightarrow \text{H}_2\text{O}(010) + \text{O}$	(t)	$2^{-11}$
37) $\text{H}_2\text{O}(020) + \text{H}_2 \rightarrow \text{H}_2\text{O}(010) + \text{H}_2$	(t)	$2^{-11}$
38) $\text{H}_2\text{O}(020) + \text{H}_2\text{O} \rightarrow \text{H}_2\text{O}(010) + \text{H}_2\text{O}$	(t)	$2^{-11}$

\* Note that the reverse rate coefficient can be calculated directly from detailed balance.

39) $\text{H}_2\text{O}(001) + \text{H} \rightarrow \text{H}_2\text{O}(000) + \text{H}$	(e)	$1^{-11}$
40) $\text{H}_2\text{O}(001) + \text{O} \rightarrow \text{H}_2\text{O}(000) + \text{O}$	(e)	$1^{-11}$
41) $\text{H}_2\text{O}(001) + \text{H}_2 \rightarrow \text{H}_2\text{O}(000) + \text{H}_2$	(o)	$2^{-13}$
42) $\text{H}_2\text{O}(001) + \text{H}_2\text{O} \rightarrow \text{H}_2\text{O}(000) + \text{H}_2\text{O}$	(q)	$1^{-12}$
43) $\text{H}_2\text{O}(001) + \text{H} \rightarrow \text{H}_2\text{O}(020) + \text{H}$	(e)	$1^{-11}$
44) $\text{H}_2\text{O}(001) + \text{O} \rightarrow \text{H}_2\text{O}(020) + \text{O}$	(e)	$1^{-11}$
45) $\text{H}_2\text{O}(001) + \text{H}_2 \rightarrow \text{H}_2\text{O}(020) + \text{H}_2$	(e)	$1^{-11}$
46) $\text{H}_2\text{O}(001) + \text{H}_2\text{O} \rightarrow \text{H}_2\text{O}(020) + \text{H}_2\text{O}$	(e)	$1^{-11}$

#### Vibration-Vibration Processes

47) $\text{H}_2(v=1) + \text{H}_2\text{O}(000)$ $\rightarrow \text{H}_2(v=0) + \text{H}_2\text{O}(001)$	(r)	$2.5^{-14} T^{5/6} \exp(-30.1/T^{1/3})$
48) $\text{H}_2(v=1) + \text{OH}(v=0)$ $\rightarrow \text{H}_2(v=0) + \text{OH}(v=1)$	(e)	$1^{-12}$
49) $\text{N}_2(v=1) + \text{H}_2\text{O}(010)$ $\rightarrow \text{N}_2(v=0) + \text{H}_2\text{O}(001)$	(e)	$3^{-13}$
50) $\text{N}_2(v=1) + \text{H}_2\text{O}(000)$ $\rightarrow \text{N}_2(v=0) + \text{H}_2\text{O}(010)$	(s)	$9.4^{-11} \exp(-64/T^{1/3})$
51) $\text{H}_2\text{O}(010) + \text{H}_2\text{O}(010)$ $\rightarrow \text{H}_2\text{O}(020) + \text{H}_2\text{O}(000)$	(e)	$1^{-12} \sqrt{T}$
52) $\text{H}_2\text{O}(001)$ and $\text{H}_2\text{O}(100)$ assumed in quasi-equilibrium at the local translational temperature.		

References (Apollo Model)

- a) D.L. Baulch, et. al., High Temperature Reaction Rate Data, Leeds University, (1969).
- b) F.P. Boynton, Semi Annual Technical Report No. 1, Physical Dynamics Report PD-72-020, February(1972).
- c) calculated from the forward rate coefficient and detailed balance, using the equilibrium constant as given in the Leeds review [(a) above].
- d) estimated from the Leeds review [(a) above] using a method outlined in (b) above.
- e) estimate
- f) W.E. Wilson Jr., J. Phys. Chem. Refer. Data 1, 535 (1972).
- g) R.F. Hiedner and V.V. Casper, Chem. Phys. Letters 15, 179(1972)
- h) J.H. Kieffer and R.W. Lutz, J. Chem. Phys. 44, 658(1966) 45, 3888(1966).
- i) C. Joffrin, J. Ducuing and J.P. Coffinet, Opt. Commun. 2, 245(1970).
- j) past measurements involving water presumably measure both VV and VT processes.
- k) W.D. Breshears and P.F. Bird, J. Chem. Phys. 48, 4768 (1968).
- l) R.J. McNeal, M.E. Whitson Jr. and G.R. Cook, preprint of work submitted to Chem. Phys. Letters(1972). Aerospace Report SPL 3303, July(1972).
- m) D.J. Eckstrom, Stanford Research Institute, unpublished results.
- n) D.R. White, J. Chem. Phys. 46, 2016(1967).
- o) estimate based on Marriott's cross section values see E.R. Fisher, AFCRL Special Report, 72-0539, August(1972).
- p) estimate based on all available data
- q) estimate based on analogy to HF VT relaxation

- r) estimate based on the modified Rapp model, J. Chem. Phys. 40, 573(1964); Fisher unpublished results.
- s) estimate based on data of C.W. Von Rosenberg, K.N.C. Bray and N.H. Pratt, J. Chem. Phys. 56, 3230(1972) and 13th Symp. (Intern.) on combustion p. 89, (1971).
- t) scaled from the analogous rate constant for exciting the lowest bending mode.
- u) estimated based on the rate for  $\text{CO} + \text{H}$ ; see reference (o) under RELAXATION MODEL (2).

TABLE II  
VIBRATIONAL RELAXATION MODEL

<u>Vibration-Translation Processes</u>	<u>Reference</u>	<u>Rate Constant (cc/molecule-sec)</u>
1) $\text{H}_2(v=1) + \text{H} \rightarrow \text{H}_2 + \text{H}$	(a)	$3.2^{-11} \exp(-1400/T)$
2) $\text{H}_2(v=1) + \text{O} \rightarrow \text{H}_2 + \text{O}$	(b)	$3.0^{-11} \sqrt{T}$
3) $\text{H}_2(v=1) + \text{H}_2 \rightarrow \text{H}_2 + \text{H}_2$	(c,d)	$1.5^{-7} \exp(-139/T^{1/3})$
4) $\text{H}_2(v=1) + \text{H}_2\text{O} \rightarrow \text{H}_2 + \text{H}_2\text{O}$	(b,e)	$1.5^{-7} \exp(-139/T^{1/3})$
5) $\text{N}_2(v=1) + \text{H} \rightarrow \text{N}_2 + \text{H}$	(y)	$1.9^{-6} \exp(-164/T^{1/3})$
6) $\text{N}_2(v=1) + \text{O} \rightarrow \text{N}_2 + \text{O}$	(f,g,h)	$1.2^{-13} \exp(-23/T^{1/3})$
7) $\text{N}_2(v=1) + \text{H}_2 \rightarrow \text{N}_2 + \text{H}_2$	(i)	$3.9^{-8} \exp(-164/T^{1/3})$
8) $\text{N}_2(v=1) + \text{H}_2\text{O} \rightarrow \text{N}_2 + \text{H}_2\text{O}$	(b,e)	$3.9^{-8} \exp(-164/T^{1/3})$
9) $\text{OH}(v=1) + \text{H} \rightarrow \text{OH} + \text{H}$	(b)	$3^{-11}$
10) $\text{OH}(v=1) + \text{O} \rightarrow \text{OH} + \text{O}$	(b)	$3^{-11}$
11) $\text{OH}(v=1) + \text{H}_2 \rightarrow \text{OH} + \text{H}_2$	(b)	$1^{-11}$
12) $\text{OH}(v=1) + \text{H}_2\text{O} \rightarrow \text{OH} + \text{H}_2\text{O}$	(b)	$3^{-12}$
13) $\text{H}_2\text{O}(010) + \text{H} \rightarrow \text{H}_2\text{O}(000) + \text{H}$	(b)	$1^{-11}$
14) $\text{H}_2\text{O}(010) + \text{O} \rightarrow \text{H}_2\text{O}(000) + \text{O}$	(b)	$1^{-11}$
15) $\text{H}_2\text{O}(010) + \text{H}_2 \rightarrow \text{H}_2\text{O}(000) + \text{H}_2$	(b,j)	$1^{-11}$
16) $\text{H}_2\text{O}(010) + \text{H}_2\text{O} \rightarrow \text{H}_2\text{O}(000) + \text{H}_2\text{O}$	(k)	$1^{-11}$
17) $\text{H}_2\text{O}(001) + \text{H} \rightarrow \text{H}_2\text{O}(000) + \text{H}$	(b)	$1^{-11}$
18) $\text{H}_2\text{O}(001) + \text{O} \rightarrow \text{H}_2\text{O}(000) + \text{O}$	(b)	$1^{-11}$
19) $\text{H}_2\text{O}(001) + \text{H}_2 \rightarrow \text{H}_2\text{O}(000) + \text{H}_2$	(j)	$2^{-13}$
20) $\text{H}_2\text{O}(001) + \text{H}_2\text{O} \rightarrow \text{H}_2\text{O}(000) + \text{H}_2\text{O}$	(l)	$1^{-12}$

21)	$\text{H}_2\text{O}(001) + \text{H} \rightarrow \text{H}_2\text{O}(020) + \text{H}$	(b)	$1^{-11}$
22)	$\text{H}_2\text{O}(001) + \text{O} \rightarrow \text{H}_2\text{O}(020) + \text{O}$	(b)	$1^{-11}$
23)	$\text{H}_2\text{O}(001) + \text{H}_2 \rightarrow \text{H}_2\text{O}(020) + \text{H}_2$	(b)	$1^{-11}$
24)	$\text{H}_2\text{O}(001) + \text{H}_2\text{O} \rightarrow \text{H}_2\text{O}(020) + \text{H}_2\text{O}$	(b)	$1^{-11}$
25)	$\text{CO}(v=1) + \text{H} \rightarrow \text{CO}(v=0) + \text{H}$	(o)	$5.3^{-7} \exp(-119/T^{1/3})$
26)	$\text{CO}(v=1) + \text{O} \rightarrow \text{CO}(v=0) + \text{O}$	(p)	$1^{-8} \exp(-96/T^{1/3})$
27)	$\text{CO}(v=1) + \text{H}_2 \rightarrow \text{CO}(v=0) + \text{H}_2$	(q)	$9^{-9} \exp(-119/T^{1/3})$
28)	$\text{CO}(v=1) + \text{H}_2\text{O} \rightarrow \text{CO}(v=0) + \text{H}_2\text{O}$	(b)	$9^{-9} \exp(-119/T^{1/3})$
29)	$\text{CO}_2(010) + \text{H} \rightarrow \text{CO}_2(000) + \text{H}$	(b)	$3.8^{-12} \exp(-7/T^{1/3})$
30)	$\text{CO}_2(010) + \text{O} \rightarrow \text{CO}_2(000) + \text{O}$	(p)	$2.3^{-9} \exp(-77/T^{1/3})$
31)	$\text{CO}_2(010) + \text{H}_2 \rightarrow \text{CO}_2(000) + \text{H}_2$	(r)	$7.6^{-12} \exp(-7/T^{1/3})$
32)	$\text{CO}_2(010) + \text{H}_2\text{O} \rightarrow \text{CO}_2(000) + \text{H}_2\text{O}$	(s)	$1^{-11}$
33)	$\text{CO}_2(010) + \text{CO}_2 \rightarrow \text{CO}_2(000) + \text{CO}_2$	(r)	$4.6^{-10} \exp(-77/T^{1/3})$
34)	$\text{CO}_2(001) + \text{H} \rightarrow \text{CO}_2(000) + \text{H}$	(t)	$6.7^{-8} \exp(-208/T^{1/3})$
35)	$\text{CO}_2(001) + \text{O} \rightarrow \text{CO}_2(000) + \text{O}$	(t)	$1^{-8} \exp(-96/T^{1/3})$
36)	$\text{CO}_2(001) + \text{H}_2 \rightarrow \text{CO}_2(000) + \text{H}_2$	(t)	$9^{-9} \exp(-119/T^{1/3})$
37)	$\text{CO}_2(001) + \text{H}_2\text{O} \rightarrow \text{CO}_2(000) + \text{H}_2\text{O}$	(t)	$9^{-9} \exp(-119/T^{1/3})$
38)	$\text{CO}_2(001) + \text{H} \rightarrow \text{CO}_2(030) + \text{H}$	(b)	$2.13^{17} T^{-6.34} \exp(-3013/T$ $-378.7/T^{1/3} + 1416/T^{2/3})$
39)	$\text{CO}_2(001) + \text{O} \rightarrow \text{CO}_2(030) + \text{O}$	(u)	$4.6^{24} T^{-5.89} \exp(-4223/T$ $-672.7/T^{1/3} + 2683/T^{2/3})$
40)	$\text{CO}_2(001) + \text{H}_2 \rightarrow \text{CO}_2(030) + \text{H}_2$	(v)	$4.27^{17} T^{-6.34} \exp(-3013/T$ $-378.7/T^{1/3} + 1416/T^{2/3})$

- 41)  $\text{CO}_2(001) + \text{H}_2\text{O} \rightarrow \text{CO}_2(030) + \text{H}_2\text{O}$  (v)  $4.71^{-40} T^{4.54} \exp(2081/T + 454/T^{1/3} - 1729/T^{2/3})$
- 42)  $\text{CO}_2(001) + \text{CO}_2 \rightarrow \text{CO}_2(030) + \text{CO}_2$  (v)  $9.16^{23} T^{-5.89} \exp(-4223/T - 672.7/T^{1/3} + 2683/T^{2/3})$
- 43)  $\text{CO}_2(001) + \text{CO} \rightarrow \text{CO}_2(030) + \text{CO}$  (v)  $6.87^{23} T^{-5.89} \exp(-4223/T - 672.7/T^{1/3} + 2683/T^{2/3})$
- 44)  $\text{CO}_2(001) + \text{N}_2 \rightarrow \text{CO}_2(030) + \text{N}_2$  (v)  $6.87^{23} T^{-5.89} \exp(-4223/T - 672.7/T^{1/3} + 2683/T^{2/3})$

#### Vibration-Vibration Processes

- 45)  $\text{H}_2(v=1) + \text{H}_2\text{O}(000)$   
 $\rightarrow \text{H}_2(v=0) + \text{H}_2\text{O}(001)$  (m)  $2.5^{-14} T^{5/6} \exp(-30.1/T^{1/3})$
- 46)  $\text{H}_2(v=1) + \text{OH}(v=0)$   
 $\rightarrow \text{H}_2(v=0) + \text{OH}(v=1)$  (b)  $1^{-12}$
- 47)  $\text{N}_2(v=1) + \text{H}_2\text{O}(010)$   
 $\rightarrow \text{N}_2(v=0) + \text{H}_2\text{O}(001)$  (b)  $3^{-13}$
- 48)  $\text{N}_2(v=1) + \text{H}_2(000)$   
 $\rightarrow \text{N}_2(v=0) + \text{H}_2\text{O}(010)$  (n)  $9.4^{-11} \exp(-64/T^{1/3})$
- 49)  $\text{H}_2\text{O}(010) + \text{H}_2\text{O}(010)$   
 $\rightarrow \text{H}_2\text{O}(020) + \text{H}_2\text{O}(000)$  (b)  $1^{-12} \sqrt{T}$
- 50)  $\text{CO}_2(001) + \text{H}_2\text{O}(000)$   
 $\rightarrow \text{CO}_2(010) + \text{H}_2\text{O}(010)$  (b)  $1^{-13}$

- 51)  $N_2(v=1) + CO_2(000)$   
 $\rightarrow N_2(v=0) + CO_2(001)$  (r) Take the largest value  $1.71^{-6} \exp(-175.3/T^{1/3})$   
 $6.07^{-14} \exp(15.3/T^{1/3})$
- 52)  $N_2(v=1) + CO(v=0)$   
 $\rightarrow N_2(v=0) + CO(v=1)$  (v) Take the largest value  $1.78^{-6} \exp(-210/T^{1/3})$   
 $6.98^{-13} \exp(-25.6/T^{1/3})$
- 53)  $CO_2(001) + CO(v=0)$   
 $\rightarrow CO_2(000) + CO(v=1)$  (b)  $1.56^{-11} \exp(-30.1/T^{1/3})$
- 54)  $CO_2(101) + H_2O(000)$   
 $\rightarrow CO_2(000) + H_2O(001)$  (b)  $2^{-13} \sqrt{T}$
- 55)  $CO_2(021) + H_2O(000)$   
 $\rightarrow CO_2(000) + H_2O(001)$  (b)  $2^{-13} \sqrt{T}$
- 56)  $CO_2(101) + CO_2(000)$   
 $\rightarrow CO_2(100) + CO_2(001)$  (b)  $2^{-11} \sqrt{T}$
- 57)  $CO_2(020) + CO_2(000)$   
 $\rightarrow CO_2(010) + CO_2(010)$  (b)  $4^{-13} \sqrt{T}$
- 58)  $H_2O(020) + H \rightarrow H_2O(010) + H$  (w)  $2^{-11}$
- 59)  $H_2O(020) + O \rightarrow H_2O(010) + O$  (w)  $2^{-11}$
- 60)  $H_2O(020) + H_2 \rightarrow H_2O(010) + H_2$  (w)  $2^{-11}$
- 61)  $H_2O(020) + H_2O \rightarrow H_2O(010) + H_2O$  (w)  $2^{-11}$
- 62)  $CO_2(020) + H \rightarrow CO_2(010) + H$  (w)  $7.6^{-12} \exp(-7/T^{1/3})$
- 63)  $CO_2(020) + O \rightarrow CO_2(010) + O$  (w)  $4.6^{-9} \exp(-77/T^{1/3})$
- 64)  $CO_2(020) + H_2 \rightarrow CO_2(010) + H_2$  (w)  $1.5^{-11} \exp(-7/T^{1/3})$
- 65)  $CO_2(020) + H_2O \rightarrow CO_2(010) + H_2O$  (w)  $2^{-11}$

- 66)  $\text{CO}_2(020) + \text{CO}_2 \rightarrow \text{CO}_2(010) + \text{CO}_2$  (w)  $9.2^{-10} \exp(-77/T^{1/3})$
- 67)  $\text{CO}_2(030) + \text{H} \rightarrow \text{CO}_2(020) + \text{H}$  (w)  $1.2^{-11} \exp(-7/T^{1/3})$
- 68)  $\text{CO}_2(030) + \text{O} \rightarrow \text{CO}_2(020) + \text{O}$  (w)  $6.9^{-9} \exp(-77/T^{1/3})$
- 69)  $\text{CO}_2(030) + \text{H}_2 \rightarrow \text{CO}_2(020) + \text{H}_2$  (w)  $2.3^{-11} \exp(-7/T^{1/3})$
- 70)  $\text{CO}_2(030) + \text{H}_2\text{O} \rightarrow \text{CO}_2(020) + \text{H}_2\text{O}$  (w)  $3^{-11}$
- 71)  $\text{CO}_2(030) + \text{CO}_2 \rightarrow \text{CO}_2(020) + \text{CO}_2$  (w)  $1.4^{-9} \exp(-77/T^{1/3})$
- 72)  $\text{H}_2\text{O}(100)$  and  $\text{H}_2\text{O}(001)$  assumed in quasi-equilibrium at the local translational temperature.
- 73)  $\text{CO}_2(100) + \text{M} \rightarrow \text{CO}_2(020) + \text{M}$  (x)

References (Vibrational Relaxation Model)

- (a) R.F. Hiedner and V.V. Casper, Chem. Phys. letters 15, 179(1972).
- (b) estimate
- (c) J.H. Kieffer and R.W. Lutz, J. Chem. Phys. 44, 658 (1966); 45, 3888(1966).
- (d) C. Joffrin, J. Ducuing and J.P. Coffinet, Opt, Commun. 2, 245(1970).
- (e) past measurements involving water presumably measure both VV and VT processes.
- (f) W.D. Breshears and P.F. Bird, J. Chem. Phys. 48, 4768 (1968).
- (g) R.J. McNeal, M.E. Whitson Jr. and G.R. Cook, preprint of work submitted to Chem. Phys. Letters (1972). Aerospace Report SPL 3303, July(1972).
- (h) D.J. Eckstrom, Stanford Research Institute, unpublished results.
- (i) D.R. White, J. Chem. Phys. 46, 2016(1967).
- (j) estimate based on Marriott's cross section values see E.R. Fisher, AFCRL Special Report, 72-0539, August (1972).
- (k) estimate based on all available data
- (l) estimate based on analogy to HF VT relaxation
- (m) estimate based on the modified Rapp model, J. Chem. Phys. 40, 573(1964) Fisher unpublished results.
- (n) estimate based on data of C.W. Von Rosenberg, K.N.C. Bray and N.H. Pratt, J. Chem. Phys. 56, 3230(1972) and 13th Symp. (Intern.) on combustion p. 89, (1971).
- (o) C.N. Rosenberg, R. Taylor and D. Teare, J. Chem. Phys. 54, 1974(1971). Also assuming the  $\text{CO}_2 + \text{H}_2$  activation energy, as shown in reaction 27).
- (p) unpublished data from AVCO.

- (q) W. J. Hoche and R. C. Millikan, J. Chem. Phys. 38, 214 (1963); and R. C. Millikan, J. Chem. Phys. 38, 2855 (1963).
- (r) R. Taylor and S. Bitterman, Rev. Mod. Phys. 41, 26 (1969).
- (s) M.I. Buckwald and S. H. Bauer, J. Phys. Chem. 76, 3108 (1972).
- (t) since direct VT relaxation of  $\text{CO}_2(v_3)$  to ground state has not been unambiguously determined without competing VV processes,  $\text{CO}_2(v_3)$  was assumed to relax like  $\text{CO}(v=1)$ .
- (u) estimate based on analogy with reaction 35).
- (v) estimate based on AVCO suggestion.
- (w) scaled from the analogous rate constant for exciting the lowest bending mode.
- (x) the Fermi resonance in  $\text{CO}_2$  is assumed to couple the (100) and (020) modes with a gas kinetic efficiency independent of collision partner.
- (y) estimated based on the rate for  $\text{CO} + \text{H}$ , under (o) above.

TABLE III  
RADIATIVE LIFETIMES

<u>SPECIES</u>	<u>t(sec)</u>
OH(v=1)	0.3
H <sub>2</sub> O(100)	0.14
H <sub>2</sub> O(010)	0.045
H <sub>2</sub> O(001)	0.014
H <sub>2</sub> O(020)	0.022
CO <sub>2</sub> (010)	0.33
CO <sub>2</sub> (001)	0.0025
CO <sub>2</sub> (101)	0.091

## SUMMARY

Two plume chemistry models are presented in this report together with the best available estimate of the rate coefficients. The intent of these models is to provide baseline chemical input to plume flow field calculations to both compare different flow field programs and to provide baseline radiation predictions in an actual plume configuration. As new rate coefficient data becomes available these models will be updated to continue to provide state-of-the-art chemistry mechanisms.

## BIBLIOGRAPHY

- Baulch, D. L., et al, "High Temperature Reaction Rate Data," Leeds University, (1969).
- Boynton, F. P., "Plume Observables and Countermeasures," Semi-Annual Technical Report No. 1, Physical Dynamics Report No. PD-72-020, February (1972).
- Boynton, F. P., E. R. Fisher, R. H. Kummler and A. Thomson, "Plume Phenomenology Handbook," Physical Dynamics Report No. PD-73-035, (1973).
- Fisher, E. R., "A Review of Vibrational Cross Sections Involving  $H_2O$ ,  $CO_2$ ,  $H_2$ ,  $CO$  and  $He$ ," AFCRL Special Report 72-0539, August (1972).
- Kummler, R. H., E. R. Fisher and F. P. Boynton, "Hydrocarbon Chemistry and Chemiluminescence in High Altitude Plumes," Physical Dynamics Report No. PD-72-031, October (1972).

APPENDIX I

Attendance List to Plume Chemistry Workshop  
Held at Physical Dynamics, Inc., Berkeley, California  
January 12, 1973

Frederick P. Boynton, Physical Dynamics

Philip E. Cassady, Lockheed-Palo Alto

Paul V. Davis, IDA

Edward R. Fisher, Physical Dynamics (Chairman)

Ralph H. Kummeler, Physical Dynamics

Paul F. Lewis, AVCO EVERETT

C. Bradley Moore, University of California-Berkeley

J. Derek Teare, AVCO EVERETT

Kenneth Wilson, Lockheed-Palo Alto

Kurt L. Wray, Boston College